



**HAL**  
open science

## **Feral cats threaten the biodiversity of French Polynesia : some island-dependent patterns**

Pauline Palmas, Jean Yves Meyer, H el ene de Meringo, Guy Teatiu, Raiarii Bambridge, Sabrina Teatiu, Tohetia Timau, Nicolas Maihota, Jean-Claude Gaertner, Eric Vidal

### **► To cite this version:**

Pauline Palmas, Jean Yves Meyer, H el ene de Meringo, Guy Teatiu, Raiarii Bambridge, et al.. Feral cats threaten the biodiversity of French Polynesia : some island-dependent patterns. III International Conference "Island Biology 2019", Jul 2019, St Denis, R eunion. . hal-03204940

**HAL Id: hal-03204940**

**<https://upf.hal.science/hal-03204940v1>**

Submitted on 22 Apr 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

# Feral cats threaten the biodiversity of French Polynesia: some island-dependent patterns

Palmas P.<sup>1,2,\*</sup>, Meyer J.-Y.<sup>3</sup>, De Méringo H.<sup>4</sup>, Teatiu G.<sup>\*</sup>, Bambridge R.<sup>\*</sup>, Teatiu S.<sup>\*</sup>, Timau T.<sup>\*</sup>, Maihota N.<sup>5</sup>, Gaertner J.-C.<sup>1,5</sup> & Vidal E.<sup>2</sup>

<sup>1</sup> University of French Polynesia, UMR-241 EIO, Labex Corail, Faa'a, Tahiti, French Polynesia

<sup>2</sup> Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale (IMBE), Aix Marseille Université, CNRS, IRD, Avignon Université, Centre IRD de Nouméa, BP A5, 98848 Nouméa cedex, Nouvelle-Calédonie

<sup>3</sup> Délégation à la Recherche, Gouvernement de la Polynésie française – B.P. 20981, 98713 Papeete, Tahiti, French Polynesia

<sup>4</sup> Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale (IMBE), Aix Marseille Université, CNRS, IRD, Avignon Université, Europôle de l'Arbois, BP 80, 13545, Aix-en-Provence, France

<sup>5</sup> IRD, UMR-241 EIO, Labex Corail, Faa'a, Tahiti, French Polynesia

\* Volunteers from the islands of Tahiti, Tahuata, Ua Huka, French Polynesia

Correspondence author : [pauline.palmas@ird.fr](mailto:pauline.palmas@ird.fr)

## BACKGROUND

Invasive populations of feral cats *Felis catus* strongly threaten the native fauna on islands worldwide (Doherty *et al.* 2016, Palmas *et al.* 2017). French Polynesia, located in the South Pacific and formed by 120 tropical and subtropical islands divided in five archipelagos, harbors several threatened endemic bird species. Feral cats which were introduced by Europeans less than 200 years ago, are currently present in most habitats of many atolls and high volcanic islands, but their abundance and impacts remain largely understudied compared to other invasive mammalian predators such as rats.


The French Polynesian islands can be considered as multi-invaded systems with different assemblages of introduced rodents. The impact of feral cat populations is assumed to be higher in presence of introduced rodents and may also vary according to the rodent species. In particular, black rats *Rattus rattus*, which constitute a constant and abundant resource, can help sustain feral cat abundance at a high level, thereby exacerbating predation pressure on native wildlife via the "hyperpredation" process (Courchamp *et al.* 1999, Ringler *et al.* 2015).

## AIMS

We investigated the impacts of feral cats on biodiversity in four contrasted inhabited high volcanic islands in French Polynesia, particularly in terms of alien rat species, by studying:

- feral cat trophic ecology and preyed endemic bird species;
- abundance of feral cat populations.

## STUDY SITES



### Marquesas Islands:


Ua Huka, 83 km<sup>2</sup> (with *R. exulans*)  
Tahuata, 69 km<sup>2</sup> (with *R. rattus*, *R. exulans*)

### Society Islands:

Moorea, 135 km<sup>2</sup> (*R. rattus*, *R. exulans*, *R. norvegicus*)  
Tahiti, 1045 km<sup>2</sup> (*R. rattus*, *R. exulans*, *R. norvegicus*)

## METHODS

### Cat diet




### Cat abundance

- 1 site per island and 2 periods of survey (2018 and 2019)
- 20 baited camera traps
- 15 days of monitoring
- 2 cam-traps per theoretical home range
- both day and night
- General Index (GI) of feral cat activity by measuring the mean of camera capture events (*i.e.* cat photos) per sampling effort

Camera trap deployment design in Vaitahu valley, Tahuata island (Marquesas)

## RESULTS

### Cat diet



### Cat abundance

Our study reveals feral cat presence and both day and night activity

Unreported patterns of cat abundance; were found: higher in the Marquesas compared to the Society, and the highest in the island of Ua Huka

Archipelago	Island	Abundance (GI index) Cat detection / 100 camera trap days
Marquesas	Ua Huka	2.1
Marquesas	Tahuata	0.5
Society	Moorea	0.03
Society	Tahiti	0.00

### Impacts on endemic bird species (IUCN Red List status according to MNHN *et al.* 2015)

<p><b>PREDATED</b></p> <p><b>LC</b> <b>Kuku</b> <i>Ptilinopus dupetithouarsii</i> White-caped fruit-dove Marquesas endemic</p>	<p><b>PREDATED</b></p> <p><b>LC</b> <b>'U'upa</b> <i>Ptilinopus purpuratus</i> Grey-green fruit-dove Society endemic</p>
<p><b>?</b></p> <p><b>CR</b> <b>Pihiti</b> <i>Vini ultramarina</i> Ultramarine lorikeet Ua Huka endemic</p>	<p><b>?</b></p> <p><b>CR</b> <b>Pahi</b> <i>Todiramphus godeffroyi</i> Marquesan kingfisher Tahuata endemic</p>
<p><b>?</b></p> <p><b>EN</b> <b>Pati'oti'o</b> <i>Pomarea iphis</i> Iphis monarch Ua Huka endemic</p>	<p><b>?</b></p> <p><b>EN</b> <b>Upe</b> <i>Ducula galeata</i> Marquesas Imperial pigeon Marquesas endemic</p>

## DISCUSSION

- We showed unreported patterns of feral cats predation in islands.
- At least 2 endemic bird species (fruit doves, Columbidae) were preyed.
- Surprising relative abundances with more feral cats on islands harboring 1 or 2 species of rodent (Ua Huka with only *Rattus exulans*; Tahuata with *R. exulans* and *R. rattus*).

## CONSERVATION IMPLICATIONS

- Although free of black rats *Rattus rattus*, the island of Ua Huka urgently needs a more in-depth study in order to elaborate an efficient management plan against feral cats.
- Bird conservation in the different islands of French Polynesia requires specific studies on invasive predators before implementation of management actions.

- The GI index could provide an alternative management tool for monitoring feral cats relative abundance between sites or changes in populations with time.

*Nota-Bene: this work is still ongoing and the analysis of data from the 2019 sampling period will confirm and clarify these patterns and conclusions (particularly concerning the bird species affected by feral cat predation)*